

Compartmentalization of the Somite and Myogenesis in Chick Embryos Are Influenced by *Wnt* Expression

J. Wagner,* C. Schmidt,* W. Nikowits, Jr.,† and B. Christ*,¹

*Institute of Anatomy, University of Freiburg, P.O. Box 111, D-79001 Freiburg, Germany; and

†School of Medicine, Stanford University, Stanford, California 94305-5151

Muscles of the body and bones of the axial skeleton derive from specialized regions of somites. Somite development is influenced by adjacent structures. In particular, the dorsal neural tube and the overlying ectoderm have been shown to be necessary for the induction of myogenic precursor cells in the dermomyotome. Members of the *Wnt* family of signaling molecules, which are expressed in the dorsal neural tube and the ectoderm, are postulated to be responsible for this process. It is shown here that ectopically implanted *Wnt-1*, *-3a*, and *-4*-expressing cells alter the process of somite compartmentalization *in vivo*. An enlarged dorsal compartment results from the implantation of *Wnt*-expressing cells ventrally between the neural tube/notochord and epithelial somites, at the expense of the ventral compartment, the sclerotome. Thus, ectopic *Wnt* expression is able to override the influence of ventralizing signals arising from notochord and floor plate. This shift of the border between the two compartments was identified by an increase in the domain of *Pax-3* expression and a complete loss of *Pax-1* expression in somites close to the ectopic *Wnt* signal. The expanded expression of *MyoD* and desmin provides evidence that it is the myotome which increases as a result of *Wnt* signaling. *Paraxis* expression is also drastically amplified after implantation of *Wnt*-expressing cells indicating that *Wnts* are involved in the formation and maintenance of somite epithelium and suggesting that *Paraxis* is activated through *Wnt* signaling pathways. Taken together these results suggest that ectopic *Wnts* disturb the normal balance of signaling molecules within the somite, resulting in an enhanced recruitment of somitic cells into the myogenic lineage. © 2000 Academic Press

Key Words: chick embryo; *Wnts*; epithelialization; somite compartmentalization; myogenesis; *Paraxis*; *Pax-3*; *Pax-1*; *MyoD*.

INTRODUCTION

Skeletal muscle of the body derives from the somites that flank the neural tube and bud off in pairs from the unsegmented paraxial mesoderm in rostral to caudal progression. The newly formed somites are epithelial spheres and subsequently become patterned to form the sclerotome ventrally and the dermomyotome dorsally. The sclerotome undergoes an epitheliomesenchymal transition and gives rise to the future vertebrae and ribs. The dermomyotome remains epithelial and becomes subdivided into dermomyotome and myotome. Myogenic precursor cells detach from, and invaginate under, the epithelial dermomyotome to form the myotome (Denetclaw *et al.*, 1997). Within the

dermomyotome two different myogenic lineages arise (Ordahl and Le Douarin, 1992; Kablar *et al.*, 1999). The medial part of the dermomyotome gives rise to epaxial muscle (trunk muscle) while the lateral part gives rise to the hypaxial muscle which will constitute the muscle of the limbs and the ventrolateral body wall (for review see Christ and Ordahl, 1995; Brand-Saberi *et al.*, 1996).

The promoting effect of dorsal neural tube and ectoderm on myogenesis coincides with the expression pattern of the *Wnt* family of genes (newly named by Nusse *et al.*, 1991). In the neural tube and in the ectoderm diverse *Wnts* are expressed (Shackleford and Varmus, 1987; Wilkinson *et al.*, 1987; Roelinck and Nusse, 1991; Parr *et al.*, 1993; Hollyday *et al.*, 1995; see also Fig. 1B). *Wnt-1* and *-3a* are expressed in the dorsal neural tube, and *Wnt-4* and *-6* in the ectoderm. *Wnts* play important roles in the determination of cell fate, proliferation, and cell survival (Nusse, 1988; McMahon,

¹ To whom correspondence should be addressed. E-mail: christb@uni-freiburg.de.

1992; McMahon and McMahon, 1989; Nusse and Varmus, 1992; Dickinson *et al.*, 1995; Yamaguchi *et al.*, 1999).

The *Wnt* genes encode secreted glycoproteins that bind to the family of Frizzled-receptors (Fz), which in turn transduce the signal into the target cells (Chan *et al.*, 1992; Bhanot *et al.*, 1996; Yang-Snyder *et al.*, 1996; He *et al.*, 1997). This leads to an accumulation of unphosphorylated β -catenin, the key mediator in a very complex signaling network (Gumbiner, 1996; Willert and Nusse, 1998; Martinez Arias *et al.*, 1999; Polakis, 1999). β -Catenin subsequently enters the nucleus where it binds to TCF/LEF transcription factors which ultimately induce the downstream Wnt target genes (Huber *et al.*, 1996a; Behrens *et al.*, 1996; Molenaar *et al.*, 1996; Clevers and van de Wetering, 1997; McKendry *et al.*, 1997; Bienz, 1998). These properties make the dorsally expressed *Wnts* excellent candidates for effectors of dorsal somite development, including the initiation of somite epithelialization, dermomyotome formation, and myogenesis.

The epithelial structure of the newly formed somite, as well as that of the dermomyotome, may be influenced by *Wnts*, which have been suggested to induce an epithelial morphology via β -catenin, in its role as a cell adhesion molecule (Hinck *et al.*, 1994; Gumbiner, 1996). Somite epithelialization, and later differentiation of the dermomyotome, can be followed by the expression of *Paraxis* (Burgess *et al.*, 1996).

Wnts were shown to be required for induction of myogenesis in the paraxial mesoderm in coculture experiments (Stern *et al.*, 1995; Münsterberg *et al.*, 1995; Fan *et al.*, 1997; Reshef *et al.*, 1998). The induction of *myf-5* expression was attributed to the neural-tube-derived *Wnts*, while *MyoD* expression was induced by ectodermally derived *Wnts* (Tajbakhsh *et al.*, 1998). Capdevila *et al.* (1998) enhanced Wnt signaling by retroviral mediated overexpression in the paraxial mesoderm and observed an enlargement of the *Pax-3*-expressing, dorsal somite compartment.

In this study we investigated the effect of ectopically expressed, dorsalizing signal molecules (*Wnt1/3a/4*) on somites *in vivo*. We focused on the questions of whether the *Wnts* are able to promote epithelialization, alter somite compartmentalization, and influence myogenesis. *Wnt*-expressing cells, when ectopically implanted between neural tube/notochord and ventral somite, produced an expansion of the dorsal, epithelially structured somitic compartment in which *Paraxis*, *Pax-3*, and *MyoD* were found to be expressed. That this expansion occurred at the expense of the ventral somitic compartment was shown by a loss of the sclerotomal marker, *Pax-1*, in affected somites.

MATERIAL AND METHODS

Embryos

Fertile eggs (White Leghorn) were obtained from a local breeder and incubated at 37.8°C and 80% relative humidity for 2 days up to HH-stages 11–14 (Hamburger and Hamilton, 1951).

TABLE 1

Implantation of Different Wnt-Secreting Cells Alters Normal Gene Expression

	<i>Pax-3</i>	<i>Pax-1</i>	<i>Paraxis</i>	<i>MyoD</i>	<i>Desmin</i>
<i>Wnt-1</i>	7/9	8/8	12/16	18/23	7/9
<i>Wnt-3a</i>	8/11	7/7	8/10	9/10	5/5
<i>Wnt-4</i>	6/6	4/4	4/4	4/4	4/5

Note. No. of embryos with altered gene expression/No. of operated embryos.

Cell Lines and Culture Conditions

Wnt-expressing rat B1a fibroblasts and NIH3T3 cells were kindly provided by A. B. Lassar and A. Kispert. These cell lines were infected with a retroviral vector, as described previously (J. Kitajewski and Z. Zheng). LNCX-/Lac-Z-transfected fibroblasts were generated in the same manner and used as control cells. The cells were selected in G418 and grown under conditions as described by Münsterberg *et al.* (1995).

Operations

Cell pellets were implanted into a prepared tunnel between the axial organs and epithelial somites by using a micropipette (Fig. 1A). The operated embryos were reincubated for approximately 20 h. The number of operated embryos is shown in Table 1.

Histology

Embryos were fixed overnight in Serra's solution, dehydrated, and embedded in paraffin. Sections were cut at 8 μ m and True red staining was carried out according to Romeis (1968).


In Situ Hybridization

Embryos were washed in PBT and fixed overnight in 4% PFA/PBT at 4°C. The antisense RNA probe was labeled with digoxigenin and whole-mount *in situ* hybridization was performed as described by Nieto *et al.* (1996). The following probes were used in this study: a probe of *Pax-1* (a 1.5-kb insert cloned into pBluescript II KS (C. Ebensperger)); a probe of *Pax-3* (a 1543-bp insert cloned into pGEM 72f (gift from M. Bronner-Fraser)); a probe of *Paraxis* (a 1380-bp insert cloned into pBSSK+ (gift from E. Olson)); and a probe of *MyoD* (a 1518-bp insert cloned into pKS+ Stratagene (Lin *et al.*, 1989)). The embryos were cryosectioned for further examinations.


RESULTS

The Influence of Wnt-Expressing Cells on Somite Compartmentalization


Wnt-1-expressing cells implanted between axial organs and epithelial somites of chick embryos at stage 12, as




Pax-3




Pax-1



Paraxis




MyoD



Desmin

Controls



shown in Fig. 1A, lead after a reincubation period of approximately 20 h to an enlarged myotome that was observed at the operation site but not at the same position in the contralateral control somites (Fig. 1C). The implanted *Wnt-1*-expressing cells are enclosed by myotomal tissue (Fig. 1D) and only single sclerotomal cells could be detected. The dermatome was not altered structurally.

***Pax-3* Expression Is Up-Regulated by *Wnt*-Expressing Cells**

The implantation of *Wnt-1*-, *Wnt-3a*-, or *Wnt-4*-expressing fibroblasts leads to increased *Pax-3* expression in adjacent somites. The grafted cells are medially and laterally flanked by a domain of strong *Pax-3* expression at the

FIG. 1. (A) Operation scheme: *Wnt*-expressing cells are grafted between axial organs and ventral epithelial somite; implanted *Wnt* cells, red; neural tube/notochord, yellow; somites, brown. (B) Control embryo (HH-stage 18) with expression patterns of the diverse *Wnts*. Data collected from Parr *et al.* (1993), Hollyday *et al.* (1995), Marcelle *et al.* (1997), and Borycki *et al.* (1998). (C) Transverse section of a HH-stage 18 embryo shows an enlarged myotome-like compartment which surrounds implanted *Wnt-1*-expressing cells. True red staining: (a) mesenchyme; (b) epithelium (D) Same embryo as shown in (C); operation side; higher magnification. (a) epithelium. In C and D, implanted cells are marked by asterisks (cross-sections). NT, neural tube; CD, notochord.

FIG. 2. *Pax-3* *in situ* hybridization. Whole-mount normal embryo; HH-stage 19. HH-stage 19 chick embryo after implantation of *Wnt-1*-expressing cells; somites show structure abnormalities. *Pax-3* is expressed throughout the whole somite and at higher levels medially and laterally. Corresponding sections of the embryo shown in (B) present an irregularly expanded expression domain; the cells are surrounded by *Pax-3*-positive tissue which is morphologically epithelially structured. Only the space between implant and notochord is left free. HH-stage 19 embryo after implantation of *Wnt-3a*-expressing cells; *Pax-3* expression is upregulated laterally in the somite at the transition to the upper limb. The whole mount of a HH-stage 18 chick embryo after implantation of *Wnt-4*-expressing cells displays a massive increase in *Pax-3* expression lateral of the cells. Corresponding sections of the embryo shown in (E) present the implanted *Wnt-4*-expressing cells lying quite dorsal in the somite. They are surrounded by *Pax-3*-positive tissue from each side. Especially strong expression is notable in the lateral part of the somite and even ventrally between cells and notochord an expression could be found. In B, D, and E, the operation site is marked by brackets (whole mounts); in C and F, implanted cells are marked by asterisks (cross-sections). NT, neural tube; CD, notochord.

FIG. 3. *Pax-1* *in situ* hybridization. (A) Whole-mount staining of a normal chick embryo HH-stage 19. (B) Chick embryo after implantation of *Wnt-1*-expressing cells shows a complete absence of *Pax-1* at the operation site. (C) Chick embryo shows an altered *Pax-1* expression after implantation of *Wnt-3a*-expressing cells. *Pax-1* is down-regulated at the operation site. (D) The whole mount of a HH-stage 19 embryo presents a complete down-regulation of *Pax-1* at the site of implantation of *Wnt-4*-expressing cells. In B–D, the operation site is marked by brackets (whole mounts).

FIG. 4. *Paraxis* *in situ* hybridization. (A) Whole-mount specimen of a normal chick embryo; HH-stage 18. (B) Corresponding transverse section. (C) Whole mount of a HH-stage 19 embryo after implantation of *Wnt-1*-expressing cells. The embryo displays an irregular somitic structure and an augmented *Paraxis* expression at the operation site. (D) Corresponding transverse section; the graft is medially surrounded by *Paraxis*-positive tissue. (E) The HH-stage 19 embryo shows an up-regulation of *Paraxis* after implantation of *Wnt-3a*-expressing cells. (F) Corresponding sections of the embryo shown in (E) display the implant lying surrounded by a tissue with a high level of *Paraxis* expression. Even the ventral domain of the somite is labeled except for the space directly neighboring the notochord. After implantation of *Wnt-4*-expressing cells the embryo displays an extreme upregulated expression of *Paraxis*. The somites are *Paraxis* positive on the whole at the operation site. Sections of the same embryo as shown in (G) present an enlarged domain of *Paraxis* expression which neighbors the implant and is epithelially structured. Ventrally a supernumerary expression domain in proximity to the notochord can be found. In C, E, and G, operation site is marked by brackets (whole mounts); in B, D, F, and H, implanted cells are marked by asterisks (cross-sections). NT, neural tube; CD, notochord.

FIG. 5. *MyoD* *in situ* hybridization After implantation of *Wnt-1*-expressing cells an altered somitic structure could be found. Sections of the same embryo as shown in (A) present the cells located between two domains of *MyoD* expression. One of them lies next to the neural tube; the other one is located ventrally in quite close proximity to the notochord. The whole mount of this embryo displays a strong up-regulation of *MyoD* at the site of implantation of *Wnt-3a*-expressing cells; HH-stage 19. (D) Sections of the same embryo as shown in (C) present an enlarged *MyoD*-positive myotome-like compartment neighboring the implanted cells. Another *MyoD*-labeled domain could be found adjacent to the cells. Embryo (HH-stage 18) after implantation of *Wnt-4*-expressing cells; graft divides the somites, which are *MyoD* positive on the whole at the operation site. The corresponding sections show three supernumerary domains of *MyoD* expression. Desmin immunohistochemistry: HH-stage 19 chick embryo after implantation of *Wnt-3a*-expressing cells; the graft is surrounded by desmin-positive cells. The muscle-forming desmin-positive compartment is enlarged and occupies even the space next to the ventral neural tube and the notochord. In A, C, and E, the operation site is marked by brackets (whole mounts); in B, D, F, and G, implanted cells are marked by asterisks (cross-sections). NT, neural tube; CD, notochord.

FIG. 6. Control cells. (A) Transverse sections show no effect on *Pax-3* expression after control cell implantation. The cells could be seen between dermomyotome and notochord surrounded by unaltered sclerotomal mesenchyme. (B) HH-stage 18 embryo after implantation of control cells; no alteration of *Pax-1* expression could be observed on the operation side. (C) Transverse section presents a normal pattern of *Paraxis* expression after implantation of control cells. The cells are located in unaltered sclerotome. (D) The transverse section shows a shortened myotome on the operation side but no extra domains of *MyoD* expression. In all panels implanted cells are marked by asterisks (cross-sections). NT, neural tube; CD, notochord.

operation site (Figs. 2B–2F). In some instances, affected somites do not show the typical gap between the medial and the lateral dermomyotomal domains of *Pax-3* expression, and thus the entire dermomyotome appears to be *Pax-3* positive. In whole-mount preparations, an increase of *Pax-3* expression could be detected especially in the lateral part of the somite at the transition to the upper limb (Fig. 2D). In transverse sections, the extended *Pax-3* domain reaches more ventrally than in unaffected somites (Fig. 2F). Only the cells in the space between the grafted cells and the notochord did not express *Pax-3* (Fig. 2C).

Pax-1 Is Down-Regulated by Wnt-Expressing Cells

After implantation of *Wnt-1*, *Wnt-3a*, or *Wnt-4*-expressing cells there was a dramatic down-regulation of *Pax-1* in the ventral part of the somites. In all cases *Pax-1* expression is clearly reduced adjacent to the implanted cells and in some embryos there was a complete absence of expression (Figs. 3B–3D).

Paraxis Expression Is Up-Regulated by Wnt-Expressing Cells

The implantation of *Wnt-1*, *Wnt-3a*, or *Wnt-4*-expressing cells leads to an increase in the domain of *Paraxis* expression (Figs. 4C–4H). Transverse sections showed that the grafted cells were surrounded by *Paraxis*-positive cells (Figs. 4D, 4F, and 4H). In some cases a ventral *Paraxis*-expressing domain was found close to the notochord (Fig. 4H).

MyoD Is Up-Regulated by Wnt-Expressing Cells

Following implantation of *Wnt-1*, *Wnt-3a*, or *Wnt-4*-expressing cells, the region of *MyoD* expression was dramatically increased (Figs. 5C–5H). In some instances, the ectopic cells divided the myotome in two parts, both of which were clearly *MyoD* positive (Fig. 5D). In other cases the implants were surrounded by an extended *MyoD*-positive myotome (Figs. 5F and 5H). *In toto* the expression domain of *MyoD* on the operated side was more than twice the width as that found on the control side.

Control Cells Do Not Alter the Expression of Pax-1, Pax-3, Paraxis, MyoD, or Desmin

To demonstrate that the up-regulation of *Pax-3*, *Paraxis*, *MyoD*, and desmin and the down-regulation of *Pax-1* was not due to a barrier effect of the implanted cells, which might inhibit normal distribution of signaling molecules arising from the neural tube/notochord, *LNCX/Lac-Z* control cells were implanted at the level of the epithelial somites between neural tube/notochord and somite. Despite a certain shortening of the dermomyotome, no alterations of the normal somite compartmentalization were found. The control cells showed no effect on the normal

expression pattern of *Pax-3* (Fig. 6A), *Pax-1* (Fig. 6B), *Paraxis* (Fig. 6C), or *MyoD* (Fig. 6D).

DISCUSSION

The present study demonstrates that ectopically implanted *Wnt-1*, *-3a*, or *-4*-expressing cells alter normal somite compartmentalization. The dorsal compartment became enlarged at the expense of the ventral compartment, as shown by the expansion of the dorsal marker, *Pax-3*, and the loss of *Pax-1* expression. The enhanced expression of *Paraxis* in the paraxial mesoderm points to an enlargement of somitic epithelium and of the dermomyotome. This enlarged compartment produced increased numbers of myogenic cells leading to an expansion of the myotome.

Somite Compartmentalization and Myogenesis Are Influenced by Wnts

Previous studies indicate that one role of signals emerging from the dorsal neural tube is to establish dorsal/ventral polarity in the somite. While Christ *et al.* (1992) described some compartmentalization of somites after ablation of the neural tube, these somites were unable to give rise to muscle. This observation emphasizes the function of the neural tube in dorsoventral patterning of the somites and in inducing myogenesis. The surface ectoderm was able to partially rescue the loss of neural tube signals, but was unable to initiate and maintain myogenesis (Bober *et al.*, 1994a; Borman and Yorde, 1994). The neural tube and surface ectoderm are both sources of Wnts, but each expresses different members of the family. The difference in Wnts expressed by the surface ectoderm and the neural tube could account for the different effect on myogenesis evoked by these two tissues.

In our experiments, no difference on somite development was observed between neural tube- (*Wnt-1* and *-3a*) and surface ectoderm- (*Wnt-4*) derived Wnts. One explanation for the strong myogenic influence of the surface ectoderm-derived *Wnt-4* signal may lie in the level of expression produced by positioning the Wnt-secreting cells close to the ventral part of the somite. Stern *et al.* (1995) demonstrated that the dosage of *Wnt-1* plays a critical role in initiating myogenesis. It is likely that the thin ectodermal layer is not able to provide the same dose of *Wnt-4* as the graft used in our experiments and that a presumably high level of *Wnt-4* can augment neural-tube-derived Wnts to increase myogenesis to the level we observed. A second possibility is that the Wnts which we ectopically expressed in the paraxial mesoderm are able to induce other Wnts in the adjacent neural tube, which in turn signal the adjacent somites. That Wnts can induce one another was shown by Marcelle *et al.* (1997), in which *Wnt-1* and *-3a* induced the expression of *Wnt-11* in the medial part of the dermomyotome.

Capdevila *et al.* (1998) showed an altered dorsoventral patterning of the somites in response to retrovirally injected

Wnt-1 in the paraxial mesoderm. Their findings are similar to ours with regard to up-regulation of *Pax-3* and loss of *Pax-1* expression. Contradictory is the fact that they did not observe any alterations in the expression pattern of the myogenic markers, while we found strong amplification of *MyoD* expression. We suggest that this discrepancy could be due to differences in the level of *Wnt-1* produced by retroviral infection and cell transplantation. Additionally, the position of the *Wnt* signal relative to the somite seems to play a critical role. Thus, we also did not find enhanced myogenesis if we implanted the *Wnt*-secreting cells under the ectoderm (data not shown).

Wnts Influence Somite Epithelialization

Paraxis expression is a prerequisite for somite epithelialization (Burgess *et al.*, 1996; Barnes *et al.*, 1997; Susic *et al.*, 1997). Ectopically implanted *Wnt*-secreting cells (*Wnt-1*, -3a, -4) up-regulate the expression of *Paraxis* and promote somite epithelialization. Thus *Paraxis* may be a target gene of the *Wnts*. This idea was supported by findings of Susic *et al.* (1997) who found *Paraxis* expression to depend on the neural tube and the surface ectoderm. They were able experimentally to switch off *Paraxis* expression and to block dermomyotome formation. Mice lacking *Paraxis* do not form somite epithelium and show severe defects of muscle (Burgess *et al.*, 1995, 1996).

One way in which the *Wnts* could influence somite epithelialization is through their suggested involvement in cell adhesion. The *Wnt* signal transduction pathway leads to an accumulation of β -catenin (Hinck *et al.*, 1994). In its role as a cell adhesion molecule, β -catenin links the cadherins to the cytoskeleton (Wheelock and Knudsen, 1991), and the cadherins represent key molecules in the process of somite epithelialization (Duband *et al.*, 1987). Additionally, the expression of *E-cadherin* depends on the activation by the TCF/LEF complex (Shimamura *et al.*, 1994; Huber *et al.*, 1996b). Linask *et al.* (1998) described the expression of *N-cadherin* and β -catenin in the myotome. Therefore, cadherins may be involved in the formation of the epithelial dermomyotome.

Wnts Diminish the Sclerotome-Inducing Effect of Shh

In our experiments, *Wnt*-expressing cells in close contact with the notochord reduced the amount of sclerotome formed, presumably by interfering with the inductive effect of Shh. Implantation of a supernumerary notochord, or the retroviral overexpression of Shh, results in altered somite patterning. The expression of *Pax-3* is lost and the whole somite expresses *Pax-1* (Pourquié *et al.*, 1993; Brand-Saberi *et al.*, 1993; Johnson *et al.*, 1994).

Conversely, several groups have found that Shh is required for myogenesis (Münsterberg *et al.*, 1995; Stern *et al.*, 1995; Buffinger and Stockdale, 1995; Cossu *et al.*, 1996; Pownall *et al.*, 1996; Dietrich *et al.*, 1997; Maroto *et al.*,

1997; Borycki *et al.*, 1998; Duprez *et al.*, 1998; Tajbakhsh *et al.*, 1998; Cann *et al.*, 1999). Therefore, a dual function of Shh has been suggested. A low concentration of Shh far from the protein source might cause a different response of somitic cells from high concentrations in close proximity to the notochord. Borycki *et al.* (1998) have suggested that different groups of somitic cells might transduce the Shh signal in different ways. They found differences in expression of *Gli* transcriptional factors and *Patched* receptors to be responsible for the dual effect of Shh. *Gli* and *Patched* are expressed in the sclerotome under the influence of the notochord and transduce the ventralizing signal to the sclerotome. *Gli 2/4* are expressed in the dorsal somite under the influence of the ectoderm and transduce the Shh signal to the myogenic lineage. Another function of Shh was recently presented by Lee *et al.* (2000). They showed that Shh induces in the sclerotome the expression of *Sfrp2*, a known *Wnt* antagonist, which could bind *Wnt-1* and *Wnt-4*, thus opposing *Wnt*-induced myogenesis. Our findings extend this observation. The elevated *Wnt* dosage in the sclerotome from the ectopic source may overwhelm the secreted *Wnt* antagonist and initiate *Wnt* binding to their receptors on ventral somitic cells. Thus, *Wnts* are able to antagonize the notochord-derived Shh signal in a dosage-dependent manner. Additional evidence for the dosage-dependent antagonism between Shh and *Wnts* derives from the study of Fan *et al.* (1997). They cultured unsegmented paraxial mesoderm explants between aggregates of Shh- and *Wnt*-secreting cells and found the mesodermal reaction to these "neighbors" to depend on the ratio of Shh to *Wnts*.

Wnts Are Part of a Gene Activation Cascade in Myogenesis

Reshef and colleagues (1998) found an elevation in *Pax-3* expression, a marker for myogenic precursors in the dermomyotome, in somites cocultured with dorsal neural tube or surface ectoderm. Expression of *Wnt-1* and -3a in the dorsal neural tube are induced by BMP4 via the overlying ectoderm (Dickinson *et al.*, 1995; Marcelle *et al.*, 1997). In experiments described here, the implantation of *Wnt-1*, -3a-, and -4-secreting cells led to an up-regulation of *Pax-3* expression. These results suggest that *Wnts* act as upstream regulators of *Pax-3*, possibly via the activity of *Paraxis*. That *Pax-3* is at least in part regulated by *Paraxis* was recently shown by Wilson-Rawls *et al.* (1999). In *Paraxis*-null mutants, *Pax-3* expression is reduced in the first somites and lost in compartmentalized somites, especially in precursors of the nonmigratory hypaxial muscle (Burgess *et al.*, 1996; Wilson-Rawls *et al.*, 1999). The incomplete loss of *Pax-3* in the *Paraxis*-null mutant could be explained by the finding of Barber *et al.* (1999) that there exist different isoforms of *Pax-3* in the migratory and in the nonmigratory myogenic lineages which might be induced via different mechanisms.

Pax-3 plays a fundamental role in the determination of myogenic cells (Franz *et al.*, 1993; Bober *et al.*, 1994b;

Goulding *et al.*, 1994; Williams and Ordahl, 1994; Tremblay *et al.*, 1998; Henderson *et al.*, 1999). Strong evidence for the role of *Pax-3* as an upstream regulator of *MyoD* expression was presented by Tajbakhsh *et al.* (1997), Maroto *et al.* (1997), and Henderson *et al.* (1999). We found an enlarged *MyoD*-positive myotome after implantation of *Wnt-1*-, *-3a*-, and *-4*-expressing cells. Therefore, the role of Wnts in regulating myogenesis within the somite may be as an upstream regulator of *MyoD* expression, indirectly via the expression of *Pax-3*. Since neither a *Pax-3*- nor a TCF-binding site has been found in the *MyoD* locus (Kucharczuk *et al.*, 1999), it is likely that intermediate factors may be necessary to activate *MyoD* expression.

The Size of the Myotome May Be Controlled by Several Mechanisms

The enlargement of the myotome observed after ectopic implantation of Wnt-secreting cells could be due to one or more factors. First, an increase in cell proliferation or a decrease in apoptosis of myogenic precursors could lead to and expanded myotome. However, in experiments not shown, we found no indication for such a mechanism. Another source of the additional myotomal cells could be the dermomyotome itself. In addition to myogenic precursors, the dermomyotome contains cells which give rise to dermis and blood vessels. Conceivably, these nonmyogenic cells could be transformed into myogenic precursor cells by a Wnt-controlled mechanism.

Finally, the ventral compartment of the somite could be another source of the additional muscle cells. In several cell culture studies the muscle-inducing ability of diverse embryonic structures was examined separately. In accordance with our results, somites exhibited myogenesis only if cultured in contact with *Wnt*-expressing structures such as neural tube and surface ectoderm (Vivarelli and Cossu, 1986; Kenny-Mobbs and Thorogood, 1987; Buffinger and Stockdale, 1994; Stern and Hauschka, 1995; Angello *et al.*, 1997). This promoting effect on dermomyotome formation and myogenesis was mimicked by *Wnt*-expressing cells in culture experiments (Münsterberg *et al.*, 1995; Stern *et al.*, 1995; Fan *et al.*, 1997). These authors observed an increase in myogenesis if Wnts acted in concert with Shh. Our results show that the enlargement of the dorsal compartment takes place at the expense of the ventral compartment. This suggests that the ventralizing effect of Shh could be diminished by the influence of dorsalizing Wnts *in vivo*.

Spence *et al.* (1996) and Dietrich *et al.* (1997) also proposed that the dorsal neural tube and notochord have opposing functions on somite myogenesis. Spence *et al.* (1996) showed that rotation of the neural tube leads to the formation of a secondary dermomyotome in the ventral somite. At the time of cell grafting, in experiments presented here, cells in the ventral part of the somite are not committed to the sclerotomal lineage (Aoyama and Asamoto, 1988; Dockter and Ordahl, 1998). Our results are

consistent with a model in which cells of the ventral somite are able to switch to a myogenic fate in response to Wnt signals emanating from the dorsal neural tube.

CONCLUSION

1. Somitic epithelium is likely induced and maintained by Wnts and might promote myogenesis.
2. Augmentation of the Wnt signal leads to an enlarged myotome.
3. Ectopically secreted Wnts diminish the sclerotome-inducing effect of Shh.
4. Wnts represent important signaling molecules for myogenic induction.
5. Wnts promote expression of *Paraxis*, *Pax-3*, and *MyoD*.

ACKNOWLEDGMENTS

We thank E. Gimbel, U. Pein, M. Schütthoff, and L. Koschny for excellent technical assistance. We are grateful to A. Lassar and A. Kispert for the gift of *Wnt*-expressing cells. The probes were kindly provided by E. Olson, M. Bronner-Fraser, and C. Ebensperger. We thank F. E. Stockdale and B. Stockdale for their constructive comments on the manuscript. This work was supported by grants from the Deutsche Forschungsgemeinschaft (Ch 44/12-3 and Ch 44/14-1) to B.C.

REFERENCES

- Angello, J., Stern, H., and Hauschka, S. (1997). P19 embryonal carcinoma cells: A model system for studying neural tube induction of skeletal myogenesis. *Dev. Biol.* **192**, 93–98.
- Aoyama, H., and Asamoto, K. (1988). Determination of somite cells: Independence of cell differentiation and morphogenesis. *Development* **104**, 15–28.
- Barber, T., Barber, M., Cloutier, T., and Friedman, T. (1999). Pax3 gene structure, alternative splicing and evolution. *Gene* **237**, 311–319.
- Barnes, G., Alexander, P., Hsu, C., Mariani, B., and Tuan, R. (1997). Cloning and characterization of chicken Paraxis: A regulator of paraxial mesoderm and somite formation. *Dev. Biol.* **189**, 95–111.
- Behrens, J., von Kries, J., Kuhl, M., Bruhn, L., Wedlich, D., Grosschedl, R., and Birchmeier, W. (1996). Functional interaction of beta-catenin with the transcription factor LEF-1. *Nature* **382**, 638–642.
- Bhanot, P., Brink, M., Harryman Samos, C., Hsieh, J., Wang, Y., Macke, J., Andrew, D., Nathans, J., and Nusse, R. (1996). A new member of the frizzled family of *Drosophila* functions as a Wingless receptor. *Nature* **382**, 225–230.
- Bienz, M. (1998). TCF: Transcriptional activator or repressor? *Curr. Opin. Cell Biol.* **10**, 366–372.
- Bladt, F., Riethmacher, D., Isenmann, S., Aguzzi, A., and Birchmeier, C. (1995). Essential role for the c-met receptor in the migration of myogenic precursor cells into the limb bud. *Nature* **376**, 768–771.
- Bober, E., Brand-Saberi, B., Ebensperger, C., Wilting, J., Balling, R., Paterson, B., Arnold, A., and Christ, B. (1994a). Initial steps of

- myogenesis in somites are independent of influence from axial structures. *Development* **120**, 3073–3082.
- Bober, E., Franz, T., Arnold, H., Gruss, P., and Tremblay, P. (1994b). Pax-3 is required for the development of limb muscles: A possible role for the migration of dermomyotomal muscle progenitor cells. *Development* **120**, 603–612.
- Borman, W., and Yorde, D. (1994). Barrier inhibition of a temporal neuraxial influence on early chick somite myogenesis. *Dev. Dyn.* **200**, 68–78.
- Borycki, A., Mendham, L., and Emerson, C. (1998). Control of somite patterning by Sonic hedgehog and its downstream signal response genes. *Development* **125**, 777–790.
- Brand-Saberi, B., Wilting, J., Ebensperger, C., and Christ, B. (1996). The formation of somite compartments in the avian embryo. *Int. J. Dev. Biol.* **40**, 411–420.
- Brand-Saberi, B., Ebensperger, C., Wilting, J., Balling, R., and Christ, B. (1993). The ventralizing effect of the notochord on somite differentiation in chick embryos. *Anat. Embryol.* **188**, 239–245.
- Buffinger, N., and Stockdale, F. (1995). Myogenic specification of somites is mediated by diffusible factors. *Dev. Biol.* **169**, 96–108.
- Buffinger, N., and Stockdale, F. (1994). Myogenic specification in somites: induction by axial structures. *Development* **120**, 1443–1452.
- Burgess, R., Rawls, A., Brown, D., Bradley, A., and Olson, E. (1996). Requirement of the paraxis gene for somite formation and musculoskeletal patterning. *Nature* **384**, 570–573.
- Burgess, R., Cserjesi, P., Ligon, K., and Olson, E. (1995). Paraxis: A basic helix–loop–helix protein expressed in paraxial mesoderm and developing somites. *Dev. Biol.* **168**, 296–306.
- Cann, G., Lee, J., and Stockdale, F. (1999). Sonic hedgehog enhances somite cell viability and formation of primary slow muscle fibers in avian segmented mesoderm. *Anat. Embryol.* **200**, 239–252.
- Capdevila, J., Tabin, C., and Johnson, R. (1998). Control of dorso-ventral somite patterning by Wnt-1 and β -catenin. *Dev. Biol.* **193**, 182–194.
- Chan, S., Karpf, D., Fowlkes, M., Hooks, M., Bradley, M., Vuong, V., Bambino, T., Liu, T., Arnaud, C., Strewler, G., and Nissenson, R. (1992). Two homologs of the *Drosophila* polarity gene frizzled are widely expressed in mammalian tissues. *J. Biol. Chem.* **267**, 25202–25207.
- Christ, B., and Ordahl, C. (1995). Early stages of chick somite development. *Anat. Embryol.* **191**, 381–396.
- Christ, B., Brand-Saberi, B., Grim, M., and Wilting, J. (1992). Local signaling in dermomyotomal cell type specification. *Anat. Embryol.* **186**, 505–510.
- Clevers, H., and van de Wetering, M. (1997). TCF/LEF factors earn their wings. *TIG* **13**, 485–489.
- Cossu, G., Kelly, R., Tajbakhsh, S., Di Donna, S., Vivarelli, E., and Buckingham, M. (1996). Activation of different myogenic pathways: Myf5 is induced by the neural tube and MyoD by the dorsal ectoderm in mouse paraxial mesoderm. *Development* **122**, 429–437.
- Denetclaw, W., Christ, B., and Ordahl, C. (1997). Location and growth of epaxial myotome precursor cells. *Development* **124**, 1601–1610.
- Dickinson, M., Selleck, M., McMahon, A., and Bronner-Fraser, M. (1995). Dorsalization of the neural tube by the non-neural ectoderm. *Development* **121**, 2099–2106.
- Dietrich, S., Schubert, F., and Lumsden, A. (1997). Control of dorsoventral pattern in the chick paraxial mesoderm. *Development* **124**, 3895–3908.
- Dockter, J., and Ordahl, C. (1998). Determination of sclerotome to the cartilage fate. *Development* **125**, 2113–2124.
- Duband, P., Dufour, S., Hatta, K., Takeichi, M., Edelman, G., and Thiery, P. (1987). Adhesion molecules during somitogenesis in the avian embryo. *J. Cell. Biol.* **104**, 1361–1374.
- Duprez, D., Fournier-Thibault, C., and Le Douarin, N. (1998). Sonic hedgehog induces proliferation of committed skeletal muscle cells in the chick limb. *Development* **125**, 495–505.
- Fan, C., Lee, C., and Tessier-Lavigne, M. (1997). A role for Wnt proteins in induction of dermomyotome. *Dev. Biol.* **191**, 160–165.
- Franz, T., Kothary, R., Surani, M., Halata, Z., and Grim, M. (1993). The splotch mutation interferes with muscle development in the limbs. *Anat. Embryol.* **187**, 153–160.
- Goulding, M., Lumsden, A., and Paquette, A. (1994). Regulation of Pax-3 expression in the dermomyotome and its role in muscle development. *Development* **120**, 957–971.
- Gumbiner, B. (1996). Cell adhesion: The molecular basis of tissue architecture and morphogenesis. *Cell* **84**, 345–357.
- Hamburger, V., and Hamilton, H. (1951). A series of normal stages in the development of the chick embryo. *J. Morphol.* **88**, 49–82.
- He, X., Saint-Jeannet, J., Wang, Y., Nathans, J., Dawid, I., and Varmus, H. (1997). A member of the Frizzled protein family mediating axis induction by Wnt-5a. *Science* **275**, 1652–1654.
- Henderson, D., Conway, S., and Copp, A. (1999). Rib truncations and fusions in the Sp2H mouse reveal a role for Pax-3 in specification of the ventrolateral and posterior parts of the somite. *Dev. Biol.* **209**, 143–158.
- Hinck, L., Nelson, W., and Papkoff, J. (1994). Wnt-1 modulates cell–cell adhesion in mammalian cells by stabilizing β -catenin binding to the cell adhesion protein cadherin. *J. Cell Biol.* **124**, 729–741.
- Hollyday, M., McMahon, J., and McMahon, A. (1995). Wnt expression patterns in chick embryo nervous system. *Mech. Dev.* **52**, 9–25.
- Huber, O., Korn, R., McLaughlin, J., Ohsugi, M., Herrmann, B., and Kemler, R. (1996a). Nuclear localization of beta-catenin by interaction with transcription factor LEF-1. *Mech. Dev.* **59**, 3–10.
- Huber, O., Bierkamp, C., and Kemler, R. (1996b). Cadherins and catenins in development. *Curr. Opin. Cell. Biol.* **8**, 685–691.
- Johnson, R., Laufer, E., Riddle, R., and Tabin, C. (1994). Ectopic expression of Sonic hedgehog alters dorsal-ventral patterning of somites. *Cell* **79**, 1165–1173.
- Kablar, B., Krastel, K., Ying, C., Tapscott, S., Goldhamer, D., and Rudnicki, M. (1999). Myogenic determination occurs independently in somites and limb buds. *Dev. Biol.* **206**, 219–231.
- Kenny-Mobbs, T., and Thorogood, P. (1987). Autonomy of differentiation in avian brachial somites and the influence of adjacent tissues. *Development* **100**, 449–462.
- Kucharczuk, K., Love, C., Dougherty, N., and Goldhamer, D. (1999). Fine-scale mapping of the MyoD core enhancer: MyoD is regulated by distinct but overlapping mechanisms in myotomal and non-myotomal muscle lineages. *Development* **126**, 1957–1965.
- Lee, C., Buttitta, L., May, N., Kispert, A., and Fan, C. (2000). SHH-N upregulates Sfrp2 to mediate its competitive interaction with WNT1 and WNT4 in the somitic mesoderm. *Development* **127**, 109–118.
- Lin, Z., Dechesne, C., Eldridge, J., and Paterson, B. (1989). An avian muscle factor related to MyoD1 activates muscle-specific promoters in nonmuscle cells of different germ-layer origin and in BrdU-treated myoblasts. *Genes Dev.* **3**, 986–996.

- Linask, K., Ludwig, C., Han, M., Liu, X., Radice, G., and Knudsen, K. (1998). N-Cadherin/catenin-mediated morphoregulation of somite formation. *Dev. Biol.* **202**, 85–102.
- Marcelle, C., Stark, M., and Bronner-Fraser, M. (1997). Coordinate actions of BMPs, Wnts, Shh and Noggin mediate patterning of the dorsal somite. *Development* **124**, 3955–3963.
- Maroto, M., Reshef, R., Muensterberg, A., Koester, S., Goulding, M., and Lassar, A. (1997). Ectopic Pax3 activates MyoD and Myf5 expression in embryonic mesoderm and neural tissue. *Cell* **89**, 139–148.
- Martinez Arias, A., Brown, A., and Brennan, K. (1999). Wnt signaling: Pathway or network? *Curr. Opin. Genet. Dev.* **9**, 447–454.
- McKendry, R., Hsu, S., Harland, R., and Grosschedl, R. (1997). LEF-1/TCF proteins mediate Wnt-inducible transcription from the *Xenopus* nodal-related 3 promoter. *Dev. Biol.* **192**, 420–431.
- McMahon, A. (1992). The Wnt family of developmental regulators. *TIG* **8**, 236–242.
- McMahon, J., and McMahon, A. (1989). Nucleotide sequence, chromosomal localization and developmental expression of the mouse int-1-related gene. *Development* **107**, 643–650.
- Molenaar, M., van de Wetering, M., Oosterwegel, M., Peterson-Maduro, J., Godsave, S., Korinek, V., Roose, J., Destree, O., and Clevers, H. (1996). XTcf-3 transcription factor mediates b-catenin-induced axis formation in *Xenopus* embryos. *Cell* **86**, 391–399.
- Münsterberg, A., Kitajewski, J., Bumcrot, D., McMahon, A., and Lassar, A. (1995). Combinatorial signaling by Sonic Hedgehog and Wnt family members induces myogenic bHLH gene expression in the somite. *Genes Dev.* **9**, 2911–2922.
- Nieto, M., Patel, K., and Wilkinson, D. (1996). In situ hybridization analysis of chick embryos in whole mount and tissue sections. *Methods Cell Biol.* **51**, 219–235.
- Nusse, R., and Varmus, H. (1992). Wnt genes. *Cell* **69**, 1073–1087.
- Nusse, R., Brown, A., Papkoff, J., Scambler, P., Shackleford, G., McMahon, A., Moon, R., and Varmus, H. (1991). A new nomenclature for int-1 and related genes: The Wnt gene family. *Cell* **64**, 231–232.
- Nusse, R. (1988). The int genes in mammary tumorigenesis and in normal development. *TIG* **4**, 291–295.
- Ordahl, C., and Le Douarin, N. (1992). Two myogenic lineages within the developing somite. *Development* **114**, 339–353.
- Parr, B., Shea, M., Vassileva, G., and McMahon, A. (1993). Mouse Wnt genes exhibit discrete domains of expression in the early embryonic CNS and limb buds. *Development* **120**, 247–261.
- Polakis, P. (1999). The oncogenic activation of β -catenin. *Curr. Opin. Genet. Dev.* **9**, 15–21.
- Pourquié, O., Coltey, M., Teillet, M., and Le Douarin, N. (1993). Control of dorsoventral patterning of somitic derivatives by notochord and floor plate. *Proc. Natl. Acad. Sci. USA* **90**, 5242–5246.
- Pownall, M., Strunk, K., and Emerson, C. (1996). Notochord signals control the transcriptional cascade of myogenic bHLH genes in somites of quail embryos. *Development* **122**, 1475–1488.
- Reshef, R., Maroto, M., and Lassar, A. (1998). Regulation of dorsal somitic cell fates: BMPs and Noggin control the timing and pattern of myogenic regulator expression. *Genes Dev.* **12**, 290–303.
- Roelink, H., and Nusse, R. (1991). Expression of two members of the Wnt family during mouse development—Restricted temporal and spatial patterns in the developing neural tube. *Genes Dev.* **5**, 381–388.
- Romeis, B. (1968). *Mikroskopische Technik*. R. Oldenbourg Verlag, München, Wein.
- Shackleford, G., and Varmus, H. (1987). Expression of the proto-oncogene int-1 is restricted to postmeiotic male germ cells and the neural tube of mid-gestational embryos. *Cell* **50**, 89–95.
- Shimamura, K., Hirano, S., McMahon, A., and Takeichi, M. (1994). Wnt-1-dependent regulation of local E-cadherin and N-cadherin expression in embryonic mouse brain. *Development* **120**, 2225–2234.
- Sosic, D., Brand-Saberi, B., Schmidt, C., Christ, B., and Olson, E. (1997). Regulation of Paraxis expression and somite formation by ectoderm- and neural tube-derived signals. *Dev. Biol.* **185**, 229–243.
- Spence, M., Yip, J., and Erickson, C. (1996). The dorsal neural tube organizes the dermomyotome and induces axial myocytes in the avian embryo. *Development* **122**, 231–241.
- Stern, H., and Hausckha, S. (1995). Neural tube and notochord promote in vitro myogenesis in single somite explants. *Dev. Biol.* **167**, 87–103.
- Stern, H., Brown, A., and Hausckha, S. (1995). Myogenesis in paraxial mesoderm: Preferential induction by dorsal neural tube and by cells expressing Wnt-1. *Development* **121**, 3675–3686.
- Tajbakhsh, S., Borello, U., Vivarelli, E., Kelly, R., Papkoff, J., Duprez, D., Buckingham, M., and Cossu, G. (1998). Differential activation of Myf5 and MyoD by different Wnts in explants of mouse paraxial mesoderm and the later activation of myogenesis in the absence of Myf5. *Development* **125**, 4155–4162.
- Tajbakhsh, S., Rocancourt, D., Cossu, G., and Buckingham, M. (1997). Redefining the genetic hierarchies controlling skeletal myogenesis. *Science* **259**, 1450–1453.
- Tremblay, P., Dietrich, S., Mericskay, M., Schubert, F., Li, Z., and Paulin, D. (1998). A crucial role for Pax-3 in the development of the hypaxial musculature and the long-range migration of muscle precursors. *Dev. Biol.* **203**, 49–61.
- Vivarelli, E., and Cossu, G. (1986). Neural control of early myogenic differentiation in cultures of mouse somites. *Dev. Biol.* **117**, 319–325.
- Wheelock, M., and Knudsen, K. (1991). N-cadherin-associated proteins in chicken muscle. *Differentiation* **46**, 35–42.
- Wilkinson, D., Bailes, J., and McMahon, A. (1987). Expression of the proto-oncogene *int-1* is restricted to specific neural cells in the developing mouse embryo. *Cell* **50**, 79–88.
- Williams, B., and Ordahl, C. (1994). Pax-3 expression in segmental mesoderm marks early stages in myogenic cell specification. *Development* **120**, 785–796.
- Willert, K., and Nusse, R. (1998). β -Catenin: A key mediator of Wnt signaling. *Curr. Opin. Genet. Dev.* **8**, 95–102.
- Wilson-Rawls, J., Hurt, C., Parsons, S., and Rawls, A. (1999). Differential regulation of epaxial and hypaxial muscle development by paraxis. *Development* **126**, 5217–5229.
- Yamaguchi, G., Bradley, A., McMahon, A., and Jones, S. (1999). Wnt5a underlies outgrowth of multiple structures in the vertebrate embryo. *Development* **126**, 1211–1223.
- Yang-Snyder, J., Miller, J., Brown, J., Lai, C., and Moon, R. (1996). A frizzled homolog functions in a vertebrate signaling pathway. *Curr. Biol.* **6**, 1302–1306.

Received for publication January 5, 2000

Revised September 1, 2000

Accepted September 1, 2000

Published online November 2, 2000